

Hyper-atoms and the critical pair Theory

Yahya O. Hamidoune*

May 22, 2008

Abstract

We introduce the notion of a hyper-atom. One of the main results of this paper is the $\frac{2|G|}{3}$ -Theorem:

Let S be a finite generating subset of an abelian group G of order ≥ 2 . Let T be a finite subset of G such that $2 \leq |S| \leq |T|$, $S + T$ is aperiodic, $0 \in S \cap T$ and

$$\frac{2|G| + 2}{3} \geq |S + T| = |S| + |T| - 1.$$

Let H be a hyper-atom of S . Then S and T are H -quasi-periodic. Moreover $\phi(S)$ and $\phi(T)$ are arithmetic progressions with the same difference, where $\phi : G \mapsto G/H$ denotes the canonical morphism.

This result implies easily the traditional critical pair Theory and its basic stone: Kemperman's Structure Theorem.

1 Introduction

For a subset A of an abelian group G , the *period* of A is $\Pi(A) = \{x \in G : A + x = A\}$. The set A is said to be *periodic* if $\Pi(A) \neq \{0\}$. A basic tool in Additive Number Theory is the following generalization of the Cauchy-Davenport Theorem due to Kneser:

Theorem A (Kneser [16]) *Let G be an abelian group and let $A, B \subset G$ be finite subsets of G such that $A + B$ is aperiodic. Then $|A + B| \geq |A| + |B| - 1$.*

The description of the subsets A and B with $|A + B| = |A| + |B| - 1$, obtained by kemperman in [12] is a deep result in the classical critical pair Theory. Another step in this direction is proposed by Grynkiewicz in [3]. The cumulated proofs of these two results is about 80 pages. One of our aims in the present work is to present a methodology leading to new easier results and shortest proofs for the existing ones. This work is essentially self-contained. We assume only Kneser's Theorem, Proposition 2, Theorem 3 and Proposition 5. The last three results are proved in around 3 pages in [10].

*UPMC Univ Paris 06, E. Combinatoire, Case 189, 4 Place Jussieu, 75005 Paris, France, hamidoune@math.jussieu.fr

The isoperimetric method is a global approach introduced by the author, which derive additive inequalities from global properties of the fragments and atoms. The reader may refer to the recent paper [10] for an introduction to the applications of this method.

For a subset X , we put $\partial_S(X) = (X + S) \setminus X$ and $X^S = G \setminus (X + S)$.

Suppose that $|G| \geq 2k - 1$ and let $0 \in S$ be a generating subset. The k th-connectivity of S is defined as

$$\kappa_k(S) = \min\{|X + S| - |X| : \infty > |X| \geq k \text{ and } |X + S| \leq |G| - k\},$$

where $\min \emptyset = |G| - 2k + 1$.

We shall say that a subset X induces a k -separation if $|X| \geq k$ and $|X^S| \geq k$. We shall say that S is k -separable if some X induces a k -separation.

A finite subset X of G such that $|X| \geq k$, $|X^S| \geq k$ and $|\partial(X)| = \kappa_k(S)$ is called a k -fragment of S . A k -fragment with minimum cardinality is called a k -atom.

Let $0 \in S$ be a generating subset of an abelian group G . We shall say that S is a *Vosper subset* if for all $X \subset G$ with $|X| \geq 2$, we have $|X + S| \geq \min(|G| - 1, |X| + |S|)$.

A subgroup with maximal cardinality which is a 1-fragment will be called a *hyper-atom*. In Section 3, we prove the existence of hyper-atoms and obtain the following result:

Let S be a finite generating subset of an abelian group G such that $0 \in S$, $|S| \leq (|G| + 1)/2$ and $\kappa_2(S) \leq |S| - 1$. Let H be a hyper-atom of S . Then $\phi(S)$ is either an arithmetic progression or a Vosper subset, where ϕ is the canonical morphism from G onto G/H .

A set A is said to be H -quasi-periodic if there is an x such that $(A \setminus (x + H)) + H = A \setminus (x + H)$.

In Section 5, we apply the global isoperimetric methodology introduced in [10] to prove the following Vosper's type result:

Let S be a finite generating subset of an abelian group G of order ≥ 2 . Let T be a finite subset of G such that $|S| \leq |T|$, $S + T$ is aperiodic, $0 \in S \cap T$ and

$$\frac{2|G| + 2}{3} \geq |S + T| = |S| + |T| - 1.$$

Let H be a hyper-atom of S . Then S and T are H -quasi-periodic. Moreover $\phi(S)$ and $\phi(T)$ are arithmetic progressions with the same difference, where $\phi : G \mapsto G/H$ denotes the canonical morphism.

This $\frac{2|G|}{3}$ -Theorem implies easily several critical pair results. As an illustration we deduce from it new proofs of Kemperman's Structure Theorem and Lev's Theorem.

Quite likely, the methods introduced in the present work lead to descriptions for subsets A, B with $|A + B| = |A| + |B| + m$, with some small other values of $m \geq 0$. However we shall limit ourselves to the case $m = -1$ in order to illustrate the method in a relatively simple context.

2 Terminology and preliminaries

Let A and B be subsets of G . The subgroup generated by A will be denoted by $\langle A \rangle$. The *Minkowski sum* is defined as

$$A + B = \{x + y : x \in A \text{ and } y \in B\}.$$

Recall the following two results:

Lemma B (*folklore*) Let G be a finite group and let A and B be subsets such that $|A| + |B| \geq |G| + 1$. Then $A + B = G$.

Theorem C (*Scherk*) [15] Let X and Y be nonempty finite subsets of an abelian group G . If there is an element c of G such that $|X \cap (c - Y)| = 1$, then $|X + Y| \geq |X| + |Y| - 1$.

Scherk's Theorem follows easily from Kneser's Theorem.

By a *proper* subgroup of G we shall mean a subgroup of G distinct from G .

The next lemma is related to a notion introduced by Lee [13]:

Lemma D [1] Let X be a subset of G . Then $(X^S)^{-S} + S = X + S$.

Clearly $X \subset (X^S)^{-S}$. Take $x \notin X + S$. Then $x \notin X^S$ and hence $x - S \subset X^S - S$. It follows that $x \in (X^S - S) + S = G \setminus ((X^S)^{-S})$. $\square$

Throughout all this section, S denotes a finite generating subset of an abelian group G with $0 \in S$. Note that the best one can get, using the isoperimetric method, for a general subset S is obtained by decomposing modulo the subgroup generated by a translated copy of S containing 0.

The reader may find all basic facts from the isoperimetric method in the recent paper [10].

Notice that $\kappa_k(S)$ is the maximal integer j such that for every finite subset $X \subset G$ with $|X| \geq k$,

$$|X + S| \geq \min \left(|G| - k + 1, |X| + j \right). \quad (1)$$

Formulae (1) is an immediate consequence of the definitions. We shall call (1) the *isoperimetric inequality*. The reader may use the conclusion of this lemma as a definition of $\kappa_k(S)$. Since $|\partial(\{0\})| \geq \kappa_1$, we have

$$\kappa_1(S) \leq |S| - 1. \quad (2)$$

The basic intersection theorem is the following:

Theorem 1 [8, 10] Let S be generating subset of an abelian group G with $0 \in S$. Let A be a k -atom and let F be a k -fragment such that $|A \cap F| \geq k$. Then $A \subset F$. In particular, distinct k -atoms intersect in at most $k - 1$ elements.

The structure of 1-atoms is the following:

Proposition 2 [6, 5]

Let S be a generating subset of an abelian group G with $0 \in S$. Let H be a 1-atom of S with $0 \in H$. Then H is a subgroup. Moreover

$$\kappa_1(S) \geq \frac{|S|}{2}. \quad (3)$$

Proof. Take $x \in H$. Since $x \in (H+x) \cap H$ and since $H+x$ is a 1-atom, we have $H+x = H$ by Theorem 1. Therefore H is a subgroup. Since S generates G , we have $|H+S| \geq 2|H|$, and hence $\kappa_1(S) = |H+S| - |H| \geq \frac{|S+H|}{2} \geq \frac{|S|}{2}$. ■

Recently, Balandraud introduced some isoperimetric objects and proved a strong form of Kneser's Theorem using Proposition 2.

The next result is proved in [7]. The finite case is reported with almost the same proof in [9].

Theorem 3 [7, 9] Let S be a finite generating 2-separable subset of an abelian group G with $0 \in S$ and $\kappa_2(S) \leq |S| - 1$. Let H be a 2-atom with $0 \in H$. Then H is a subgroup or $|H| = 2$.

A short proof of this result is given in [10].

Corollary 4 [[7], Theorem 4.6] Let S be a 2-separable finite subset of an abelian group G such that $0 \in S$, $|S| \leq (|G| + 1)/2$ and $\kappa_2(S) \leq |S| - 1$.

If S is not an arithmetic progression then there is a subgroup H which is a 2-fragment of S .

Proof.

Suppose that S is not an arithmetic progression.

Let H be a 2-atom such that $0 \in H$. If $\kappa_2 \leq |S| - 2$, then clearly $\kappa_2 = \kappa_1$ and H is also a 1-atom. By Proposition 2, H is a subgroup. Then we may assume

$$\kappa_2(S) = |S| - 1.$$

By Theorem 3, it would be enough to consider the case $|H| = 2$, say $H = \{0, x\}$. Put $N = \langle x \rangle$.

Decompose $S = S_0 \cup \dots \cup S_j$ modulo N , where $|S_0 + H| \leq |S_1 + H| \leq \dots \leq |S_j + H|$. We have $|S| + 1 = |S + H| = \sum_{0 \leq i \leq j} |S_i + \{0, x\}|$.

Then $|S_i| = |N|$, for all $i \geq 1$. We have $j \geq 1$, since otherwise S would be an arithmetic progression. In particular N is finite. We have $|N + S| < |G|$, since otherwise $|S| \geq |G| - |N| + 1 \geq \frac{|G|+2}{2}$, a contradiction.

Now

$$\begin{aligned} |N| + |S| - 1 &= |N| + \kappa_2(S) \\ &\leq |S + N| = (j + 1)|N| \\ &\leq |S| + |N| - 1, \end{aligned}$$

and hence N is a 2-fragment. ■

Corollary 4 was used to solve Lewin's Conjecture on the Frobenius number [9]. Corollary 4 coincides with [[7], Theorem 4.6]. A special case of this result is Theorem 6.6 of [9]. As mentioned in [11], there was a misprint in this last statement. Indeed $|H| + |B| - 1$ should be replaced by $|H| + |B|$ in case (iii) of [Theorem 6.6, [9]].

Alternative proofs of Corollary 4 (with $|S| \leq |G|/2$ replacing $|S| \leq (|G| + 1)/2$), using Kemperman's Structure Theorem, were obtained by Gryniewicz in [2] and Lev in [14]. In the present paper, Corollary 4 will be one of the pieces leading to a new proof of Kemperman's Theorem.

Let H be a subgroup. A partition $A = \bigcup_{i \in I} A_i$ will be called a H -decomposition of A if for every i , where A_i is the nonempty intersection of some H -coset with A . A H -decomposition $A = \bigcup_{i \in I} A_i$ will be called a H -modular-progression if it is an arithmetic progression modulo H .

We need the following consequence of Menger's Theorem:

Proposition 5 [10] *Let G be an abelian group and let S be a finite subset of G with $0 \in S$. Let H be a subgroup of G and let $S = S_0 \cup \dots \cup S_u$ be a H -decomposition with $0 \in S_0$. Let $X = X_0 \cup \dots \cup X_t$ be a H -decomposition with $\frac{|G|}{|H|} \geq t + u + 1$. Assume that $\kappa_1(\phi(S)) \geq u$.*

Then there are pairwise distinct elements $n_1, n_2, \dots, n_r \in [0, t]$ and elements $y_1, y_2, \dots, y_r \in S \setminus H$ such that

$$|\phi(X \cup X_{n_1}y_1 \cup \dots \cup X_{n_r}y_r)| = t + u + 1.$$

We call the property given in Proposition 5 the *strong isoperimetric property*.

3 Hyper-atoms

In this section, we investigate the new notion of a hyper-atom. Recall that S is a Vosper subset if and only if S is non 2-separable or if $\kappa_2(S) \geq |S|$.

Lemma 6 *Let S be a finite generating Vosper subset of an abelian group G such that $0 \in S$. Let $X \subset G$ be such that $|X + S| = |X| + |S| - 1$. Also assume that $|X| \geq |S|$ if $|X| + |S| = |G|$. Then for every $y \in S$, we have $|X + (S \setminus \{y\})| \geq |X| + |S| - 2$.*

Proof.

By the definition of a Vosper subset, we have $|X + S| \geq |G| - 1$. There are two possibilities:

Case 1. $|X + S| = |G| - 1$.

Suppose that $|X + (S \setminus \{y\})| \leq |X| + |S| - 3$ and take an element z of $(X + S) \setminus (X + (S \setminus \{y\}))$. We have $z - y \in X$. Also $(X \setminus \{z - y\}) + S \subset ((X + S) \setminus \{z\})$. By the definition of a Vosper

subset, we have $|(X \setminus \{z - y\}) + S| \geq \min(|G| - 1, |X| - 1 + |S|) = |X| + |S| - 1$. Clearly $X + S \supset ((X \setminus \{z - y\}) + S) \cup \{z\}$. Hence $|X + S| \geq |X| + |S|$, a contradiction.

Case 2. $|X + S| = |G|$.

Suppose that $|X + (S \setminus \{y\})| \leq |X| + |S| - 3$ and take a 2-subset R of $(X + S) \setminus (X + (S \setminus \{y\}))$. We have $R - y \subset X$. Also $(X \setminus (R - y)) + S \subset (X + S) \setminus R$. By the definition of a Vosper subset, $|(X \setminus (R - y)) + S| \geq \min(|G| - 1, |X| - 2 + |S|)$. We have $|X| = 1$. Otherwise and since $X + S \supset ((X \setminus (R - y)) + S) \cup R$, we have $|X + S| \geq |X| + |S|$, a contradiction. Then $|X| = |S| = 3$, and hence $|G| = 5$. Now by the Cauchy Davenport Theorem, $|X + (S \setminus \{y\})| \geq |X| + |S| - 2$, a contradiction. ■

Let us prove a lemma about the fragments in quotient groups.

Lemma 7 *Let G be an abelian group and let S be a finite 2-separable generating subset containing 0. Let H be a subgroup which is a 2-fragment and let $\phi : G \mapsto G/H$ be the canonical morphism. Then*

$$\kappa_1(\phi(S)) = |\phi(S)| - 1. \quad (4)$$

Let K be a subgroup which is a 1-fragment of $\phi(S)$. Then $\phi^{-1}(K)$ is a 2-fragment of S .

Proof.

Put $|\phi(S)| = u + 1$. Since $|G| > |H + S|$, we have $\phi(S) \neq G/H$, and hence $\phi(S)$ is 1-separable.

Let $X \subset G/H$ be such that $X + \phi(S) \neq G/H$. Clearly $\phi^{-1}(X) + S \neq G$. Then $|\phi^{-1}(X) + S| \geq |\phi^{-1}(X)| + \kappa_2(S) = |\phi^{-1}(X)| + u|H|$.

It follows that $|X + \phi(S)||H| \geq |X||H| + u|H|$. Hence $\kappa_1(\phi(S)) \geq u = |\phi(S)| - 1$. The reverse inequality is obvious and follows by (2). This proves (4).

Let K be a subgroup which is a 1-fragment of $\phi(S)$. Then $|K + \phi(S)| = |K| + u$. Thus $|\phi^{-1}(K) + S| = |K||H| + u|H|$. In particular, $\phi^{-1}(K)$ is a 2-fragment. ■

Let S be a finite generating subset of an abelian group G such that $0 \in S$. Proposition 2 states that there is a 1-atom of S which is a subgroup. A subgroup with maximal cardinality which is a 1-fragment will be called a *hyper-atom* of S . This definition may be adapted to non-abelian groups and even abstract graphs. As we shall see, the hyper-atom is more closely related to the critical pair theory than the 2-atom.

Theorem 8 *Let S be a finite generating subset of an abelian group G such that $0 \in S$, $|S| \leq (|G| + 1)/2$ and $\kappa_2(S) \leq |S| - 1$. Let H be a hyper-atom of S . Then*

$\phi(S)$ is either an arithmetic progression or a Vosper subset, where ϕ is the canonical morphism from G onto G/H .

Proof.

Let us show that

$$2|\phi(S)| - 1 \leq \frac{|G|}{|H|}. \quad (5)$$

Clearly we may assume that G is finite.

Observe that $2|S + H| - 2|H| \leq 2|S| - 2 < |G|$. It follows, since $|S + H|$ is a multiple of $|H|$, that $2|S + H| \leq |G| + |H|$, and hence (5) holds.

Suppose now that $\phi(S)$ is not a Vosper subset. By the definition of a Vosper subset, $\phi(S)$ is 2-separable and $\kappa_2(\phi(S)) \leq |\phi(S)| - 1$.

Observe that $\phi(S)$ can not have a 1-fragment M which is a non-zero subgroup. Otherwise by Lemma 7, $\phi^{-1}(M)$ is a 2-fragment of S containing strictly H , contradicting the maximality of H . By (5) and Corollary 4, $\phi(S)$ is an arithmetic progression. ■

Theorem 8 implies a result proved by Plagne and the author [11] and some extensions of it, proved using Kermperman's Theory, obtained by Gryniewicz in [2] and Lev in [14].

All these results follow from Theorem 8. Two main new facts in Theorem 8 are:

- The subgroup H in Theorem 8 is well described as a hyper-atom;
- The equality $|H + S| - |H| = \kappa_1$ is much precise than the inequality $|H + S| \leq |H| + |S| - 1$ in the previous results. This equality will be needed later.

4 Transfer Lemmas

Lemma 9 *Let G be an abelian group and let $Y \subset G$ be an arithmetic progression containing 0. Let $X \subset \langle Y \rangle$ such that $|X + Y| = |X| + |Y| - 1$. Suppose that $|\langle Y \rangle| \neq |X + Y|$ or $|Y| = 2$. Then X is an arithmetic progression with the same difference as Y .*

Proof. The Lemma is obvious, once observed that a subset with cardinality $|\langle Y \rangle| - 1$ of the cyclic group $\langle Y \rangle$ (generated by an arithmetic progression containing 0) is an arithmetic progression with arbitrary difference. ■

Lemma 10 *Let S and T be finite subsets of an abelian group G such that $0 \in S \cap T$ and $S + T$ is aperiodic. Also assume that $2 \leq |S|$ and that $|S + T| = |S| + |T| - 1$. If $T \not\subset \langle S \rangle$ then T is M -quasi-periodic, where $\langle S \rangle = M$.*

Proof. Decompose $T = T_1 \cup \dots \cup T_t$ modulo M . Put $W = \{i \in [0, t] : |T_i + S| < |M|\}$. Therefore we have using (3),

$$\begin{aligned}
|T + S| &= \sum_{i \in W} |T_i + S| + \sum_{i \notin W} |M| \\
&\geq \sum_{i \in W} (|T_i| + \kappa_1(S)) + \sum_{i \notin W} |T_i + S| \\
&\geq |T| + |W| \frac{|S|}{2}.
\end{aligned}$$

It follows that $W = \{j\}$, for some j . Since $T + S$ is aperiodic, we have by Kneser's Theorem, $|T_j + S| \geq |T_j| + |S| - 1$. Since

$$|T| + |S| - 1 = |T + S| \geq \sum_{i \neq j} |T_i + S| + |T_j| + |S| - 1,$$

the result follows ■

Lemma 11 *Let S and T be finite subsets of an abelian group G such that $0 \in S \cap T$ and $S + T$ is aperiodic. Let H be a finite subgroup and let $\phi : G \mapsto G/H$ denotes the canonical morphism. Let $S = \bigcup_{0 \leq i \leq u} S_i$ be a H -decomposition such that $\phi(S_0), \dots, \phi(S_u)$ is a progression with difference d and that $S \setminus S_u$ is H -periodic. Then there is a decomposition $T = \bigcup_{0 \leq i \leq t} T_i$, such that $\phi(T_0), \dots, \phi(T_t)$ is a progression with difference d and such that $T \setminus T_t$ is H -periodic.*

The proof is an easy exercise.

Lemma 12 *Let S and T be subsets of a finite abelian group G , generated by S such that $S + T$ is aperiodic, $0 \in S \cap T$ and $|S + T| = |S| + |T| - 1$. Then $T^S - S$ is aperiodic and $|T^S - S| = |T^S| + |S| - 1$.*

Proof. The set $T^S - S$ is aperiodic by Lemma D. Clearly $T^S - S \subset G \setminus T$. Thus $|T^S - S| \leq |G| - |T| = |T^S| + |S| - 1$. By Kneser's Theorem we have $|T^S - S| \geq |T^S| + |S| - 1$. ■

5 The $\frac{2|G|}{3}$ -Theorem

The following result encodes efficiently the critical pair Theory.

Theorem 13 *Let S be a finite generating subset of an abelian group G of order ≥ 2 . Let T be a finite subset of G such that $|S| \leq |T|$, $S + T$ is aperiodic, $0 \in S \cap T$ and*

$$\frac{2|G| + 2}{3} \geq |S + T| = |S| + |T| - 1.$$

Let H be a hyper-atom of S and let $\phi : G \mapsto G/H$ denotes the canonical morphism. Then

- S and T are H -quasi-periodic,
- $\phi(S)$ and $\phi(T)$ are arithmetic progressions with the same difference.

Proof.

Set $|G| = n$, $h = |H|$, $|\phi(S)| = u + 1$, $|\phi(T)| = t + 1$ and $q = \frac{n}{h}$.

We have $|S| \leq \frac{|S|+|T|}{2} \leq \frac{n+1}{3} < \frac{n+1}{2}$.

Assume first that $|H| = 1$. By Corollary 4, S is an arithmetic progression. The result holds clearly, observing that S generates G .

Assume now that $|H| \geq 2$. Take H -decompositions $S = \bigcup_{0 \leq i \leq u} S_i$, $T = \bigcup_{0 \leq i \leq t} T_i$ and $S + T = \bigcup_{0 \leq i \leq k} E_i$.

Without loss of generality, we shall assume that

- $0 \in S_0$ and $|S_0| \geq \dots \geq |S_u|$.
- $|E_{t+1}| \geq \dots \geq |E_k|$ and $T_i + S_0 \subset E_i$, for all $0 \leq i \leq t$.

For $0 \leq i \leq t$, we put $\alpha_i = |E_i| - |T_i|$. By the definition we have $u|H| = |H + S| - |H| = \kappa_2(S) \leq |S| - 1$. It follows that for all $u \geq j \geq 0$

$$|S_{u-j}| + \dots + |S_u| \geq j|H| + 1 \quad (6)$$

Put $P = \{i \in [0, t] : |E_i| = h\}$, $W = \{i \in [0, t] : |E_i| < h\}$.

By (6), $|S_0| > \frac{h}{2}$. Thus $\langle S_0 \rangle = H$, by Lemma B. We have by (3), $|T_i + S_0| \geq |T_i| + \kappa_1(S_0) \geq |T_i| + \frac{|S_0|}{2}$, for all $i \in W$. Therefore

$$\begin{aligned} |T + S| &\geq \sum_{i \in W} |T_i + S_0| + \sum_{i \in P} |E_i| + \sum_{t+1 \leq i \leq k} |E_i| \\ &\geq |T| + |W| \frac{|S_0|}{2} + \sum_{i \in P} \alpha_i + \sum_{t+1 \leq i \leq k} |E_i|. \end{aligned} \quad (7)$$

Claim 0 $t + 1 + u \leq q$.

Suppose the contrary. By Lemma B, $\phi(S \setminus S_u) + \phi(T) = G/H$. In particular $k + 1 = q$ and $|E_i| \geq |S_{u-1}|$, for all i . Therefore we have by (6),

$$\begin{aligned} |S + T| &\geq (t + 1)|S_0| + (q - t - 1)|S_{u-1}| \\ &= (2t + 2 - q)|S_0| + (q - t - 1)(|S_0| + |S_{u-1}|) \\ &= (2t + 2 - q) \frac{uh + 1}{u + 1} + 2 \frac{uh + 1}{u + 1} (q - t - 1) \\ &\geq \frac{un + 3}{u + 1}, \end{aligned}$$

noticing that $q < t + 1 + u \leq 2t + 1$. Therefore $u = 1$ and $q = t + 1$.

Since $\frac{n}{3} > |S| - 1 \geq \kappa_2(S) \geq h = \frac{n}{q}$, we must have $q \geq 4$.

We must have $|W| \geq 3$, since otherwise by (7),

$$|T + S| \geq (q - 2)h + 2|S_0| \geq (q - 1)h + 1 = n - h + 1 > \frac{2n}{3} + 1,$$

a contradiction. We must have $|W| = 3$, since otherwise by (7), $|T + S| \geq |T| + 2|S_0| \geq |T| + |S|$, a contradiction. Then $|P| = q - |W| \geq 4 - 3 = 1$.

Since $\langle S \rangle = G$ and $u = 1$, we have $\langle \phi(S) \rangle = \langle \phi(S_1) \rangle = G/H$, and hence there a $\gamma \in P$ such that $T_\gamma + S_1 \subset E_j$, for some $j \in W$.

By Lemma B, $|T_\gamma| + |S_j| \leq h$. In particular, $\alpha_\gamma \geq |S_1|$. By (7), $|T + S| \geq |T| + |S_1| + \frac{3|S_0|}{2} > |T| + |S|$, a contradiction. Hence $t + 1 + u \leq q$.

Claim 1. $|\phi(S + T)| = |\phi(S)| + |\phi(T)| - 1$.

By Claim 0, (4) and (1), we have

$$k + 1 = |\phi(S + T)| \geq \min(q, t + 1 + u) = t + 1 + u.$$

Also, we have $(t + 1)h \geq |T| \geq |S| > \kappa_2(S) = uh$, and hence $t \geq u$. By Lemma 4, $\kappa_1(\phi(S)) = |\phi(S)| - 1$.

By Proposition 5, there a subset $J \subset [0, t]$ with $|J| = u$ and a family $\{m_i; i \in J\}$ of integers in $[1, u]$ such that $T + S$ contains the H -decomposition $(\bigcup_{0 \leq i \leq t} T_i + S_0) \cup (\bigcup_{i \in J} T_i + S_{m_i})$.

We have $k = t + u$, since otherwise

$$\begin{aligned} |S + T| &\geq \sum_{i \in J} |T_i + S_0| + \sum_{i \notin J} |T_i + S_0| + \sum_{i \in J} |T_i + S_{m_i}| + \sum_{i \geq t+u+1} |E_i| \\ &\geq u|S_0| + |T| + |S_u| \geq |T| + |S|, \end{aligned}$$

a contradiction. Thus

$$|\phi(S + T)| = |\phi(S)| + |\phi(T)| - 1.$$

Claim 2 $|E_{k-1}| \geq |S_{u-1}|$.

By Theorem 8, $\phi(S)$ is an arithmetic progression or a Vosper subset. Let us show that

$$|\phi(T) + \phi(S \setminus S_0)| \geq t + u \tag{8}$$

Notice that (8) is obvious if $\phi(S)$ is an arithmetic progression and follows by Lemma 6 if $\phi(S)$ is a Vosper subset. Claim 2 follows now.

Claim 3 If $u \geq 2$ then $q - 1 \geq t + u + 2$.

Assume $u \geq 2$. We must have

$$|P| \geq 2. \tag{9}$$

Suppose the contrary. By Lemma B, $|T_i| < \frac{h}{3} < \frac{|S_0|}{2}$ for every $i \in W$. We have using Claim 2 and (6),

$$\begin{aligned} 2|T| > |S + T| &\geq \sum_{i \in W} |T_i + S_0| + \sum_{i \in P} |E_i| + \sum_{t+1 \leq i \leq k} |E_i| \\ &\geq \sum_{i \in W} 2|T_i| + |P||H| + |S_{u-1}| + |S_u| \\ &\geq \sum_{i \in W} 2|T_i| + 2h + 1 > 2|T|, \end{aligned}$$

a contradiction. Take $U \subset P$ with $|U| = 2$. We have using (6),

$$2|S_0| \geq |S_0| + |S_{u-1}| \geq \frac{2}{3}(|S_u| + |S_{u-1}| + |S_{u-2}|) \geq \frac{4h+2}{3}.$$

By (8), we have $|E_i| \geq |S_{u-1}|$ for all $i \leq k-1$. Clearly $|E_i| \geq |S_0|$, for all $0 \leq i \leq t$. The claim must hold since otherwise we have using Claim 2 and (6),

$$\begin{aligned} |S+T| &\geq \sum_{i \in U} |E_i| + (t-1)|S_0| + (u-1)|S_{u-1}| + |S_u| \\ &= 2h + (t-2)|S_0| + (u-2)|S_{u-1}| + (|S_0| + |S_{u-1}| + |S_u|) \\ &= 2h + (t-u)|S_0| + (u-2)(|S_0| + |S_{u-1}|) + (|S_0| + |S_{u-1}| + |S_u|) \\ &\geq 2h + (t-u)\frac{2h+1}{3} + \frac{(4h+2)(u-2)}{3} + 2h+1 \\ &> (t+u+2)\frac{2h}{3} + 1 \geq q\frac{2h}{3} + 1 = \frac{2n}{3} + 1, \end{aligned}$$

a contradiction.

By Claim 1 and Claim 3, $\phi(S)$ can not be a Vosper subset. By Theorem 8, $\phi(S)$ is an arithmetic progression. By Lemma 8 and since $t+1+u < q$ for $u \geq 2$, $\phi(T)$ is an arithmetic progression with the same difference as $\phi(S)$.

Now we shall reorder the S_i 's And T_i 's using the modular progression structure.

Take H -decompositions $S = \bigcup_{0 \leq i \leq u} B_i$, $T = \bigcup_{0 \leq i \leq t} A_i$ and a H -decomposition $S+T = \bigcup_{0 \leq i \leq k} R_i$. Since $-d$ is a difference of $\phi(S)$, we may assume $0 \in B_0$ and that

1. $\phi(B_0), \dots, \phi(B_u)$ is an arithmetic progression with difference d and $|B_0| \geq |B_u|$.
2. $\phi(A_0), \dots, \phi(A_t)$ is an arithmetic progression with difference d .
3. $A_i + B_0 \subset R_i$, for all $0 \leq i \leq t$;
4. $A_t + B_i \subset R_{t+i}$, for all $1 \leq i \leq u$.

We shall put $Y = \{i \in [0, t] : |R_i| < h\}$. Since $|B_0| \geq |B_u|$, we have using (6), $|B_0| > \frac{h}{2}$. Thus $\langle B_0 \rangle = H$, by Lemma B. We have by (3), $|A_i + B_0| \geq |A_i| + \kappa_1(B_0) \geq |A_i| + \frac{|B_0|}{2}$, for all $i \in Y$. Thus

$$|T+S| \geq \sum_{0 \leq i \leq t} |A_i + B_0| + \sum_{1 \leq i \leq u} |A_t + B_i| \tag{10}$$

$$\geq |T| + |Y|\frac{|B_0|}{2} + \sum_{1 \leq i \leq u} |A_t + B_i|. \tag{11}$$

By (11), we have $|T+S| \geq |T| + |Y|\frac{|B_0|}{2} + |S \setminus B_0|$, and hence $|Y| \leq 1$.

Claim 4. $Y = \emptyset$. Suppose the contrary. Then $Y = \{r\}$, for some $0 \leq r \leq t$. Assume first $r < t$. By Lemma B, $h \geq |A_r| + |B_0|$. Thus

$$\begin{aligned} |S + T| &\geq |R_r| + th + |A_t + (B \setminus B_0)| \\ &\geq |B_0| + |A_r| + |B_0| + (t-1)h + |A_t| + \sum_{1 \leq i \leq u-1} |B_i| \\ &\geq |T| + |S| - |B_u| + |B_0| \geq |S| + |T|, \end{aligned}$$

a contradiction. Then $r = t$. By Lemma B, $h \geq |A_t| + |B_0|$. Also $|R_t| \geq |A_{t-1}|$. Hence

$$\begin{aligned} |S + T| &\geq th + |R_t| + |A_t + (B \setminus B_0)| \\ &\geq |A_t| + |B_0| + (t-1)h + |A_{t-1}| + \sum_{1 \leq i \leq u} |B_i| \\ &\geq |T| + |S|, \end{aligned}$$

a contradiction.

Let us show that $|E_i| = h$, for all $i \leq k-1$.

Suppose that there is an $r \leq k-1$ with $|E_r| < h$. By Claim 4, $t+1 \leq r$. Since $(A_t + B_r) \cup (A_{t-1} + B_{r+1}) \subset R_r$, we have using (6), $2h \geq |A_t| + |B_r| + |A_{t-1}| + |B_{r+1}| \geq |A_t| + |A_{t-1}| + h + 1$, by Lemma B. Thus $|T+H| - |T| \geq 2h - (|A_t| + |A_{t-1}|) \geq h + 1$. Now $|S+T| \geq |T+H| + |S \setminus B_0| \geq |T| + h + 1 + |S| - |B_0| > |T| + |S|$, a contradiction.

Since $S + T$ is aperiodic, the set $B_u + A_t$ is aperiodic. By Kneser's Theorem $|B_u + A_t| \geq |B_u| + |A_t| - 1$. Now have

$$\begin{aligned} |S| + |T| - 1 = |S + T| &\geq (t+u)h + |A_t + B_u| \\ &\geq th + |B_u| + th + |A_t| - 1 + \sum_{1 \leq i \leq u-1} |B_i| \\ &\geq |T| + |S| - 1 \end{aligned}$$

Thus $|S \setminus B_u| = uh$ and $|T \setminus A_t| = th$. ■

Notice that the subgroup in Theorem 13 depends only one of the sets (namely S), while the subgroup in Kemperman Structure Theorem depends on S and T .

6 The Structure Theory

A pair $\{A, B\}$ of subsets of an abelian group G will be called a *weak pair* if one of the following conditions holds:

- (WP1) $\min(|A|, |B|) = 1$;
- (WP2) There is $d \in G$, with order $\geq |A| + |B| - 1$, such that A and B are arithmetic progressions with difference d ;
- (WP3) A is aperiodic and there is a finite subgroup H and $g \in G$ such that A, B are contained in some H -cosets and $g - B = H \setminus A$;

(WP4) There is a subgroup H such that A, B are contained in some H -cosets and $|A| + |B| = |H| + 1$, and moreover there is $c \in G$ such that $|(c - A) \cap B| = 1$.

An elementary pair is a pair $\{A, B\}$ satisfying one of the conditions (WP1), (WP2), (SP3) and (SP4), where

- $SP3 = WP3$ and for every $c \in G$, such that $|(c - A) \cap B| \neq 1$,
- $SP4 = WP4$ and for every $d \in G \setminus \{c\}$, $|(d - A) \cap B| \neq 1$.

The notion of a weak pair was suggested by Kemperman [12] (end of section 5, Page 82) in order to formulate an easier description. The reader could use the following Lemma, implicit in Kemperman's work [12], if he wants to work with elementary pairs.

Lemma E (*Kemperman*) *Let A, B be a weak pair.*

There is a nonzero subgroup H and H -quasi-periodic decompositions $A = A_0 \cup A_1$ and $B = B_0 \cup B_1$ such that (A_1, B_1) is an elementary pair and $|(\phi(a_1 + b_1) - \phi(A)) \cap \phi(B)| = 1$, where $\phi : G \mapsto G/H$ is the canonical morphism and $a_1 \in A_1$ and $b_1 \in B_1$.

The proof of the above lemma is implicit Kemperman's work [12]. It can be done within one page or less using Kneser's Theorem. The reader may also refer to the Appendix of Lev's paper [14].

Theorem 14 *Let T and S be a finite subset of an abelian G such that $\langle S \cup T \rangle = G$. Assume that $|S| \leq |T|$, $0 \in S \cap T$ and*

$$|S + T| = |S| + |T| - 1.$$

Also assume that $S + T$ is aperiodic and that $\{S, T\}$ is not a weak pair.

- (i) *If $G \neq \langle S \rangle$ then T is $\langle S \rangle$ -quasi-periodic;*
- (ii) *If $G = \langle S \rangle$ and $G \neq \langle U \rangle$ then S and T are U -quasi-periodic, where $U = T^S - a$, for some $a \in T^S$;*
- (iii) *If $G = \langle S \rangle = \langle U \rangle$ then there is a proper subgroup H of L such that S and T are H -quasi-periodic modular progressions. Moreover H is either a hyper-atom of S or a hyper-atom of U .*

Proof. Since $\{S, T\}$ is not a weak pair, we have $|S| \geq 2$. Put $L = \langle S \rangle$. If $L \neq G$, then T is L -quasi-periodic by Lemma 10 and (i) holds. So we may assume without loss of generality that S generates G . We have $|T^S| \geq 2$, otherwise putting $G = (T + S) \cup \{v\}$, we have clearly $v - S \subset G \setminus T$. Actually we have equality by the relation $|T + S| = |T| + |S| - 1$. Then $\{T, S\}$ is a weak pair, a contradiction.

Notice that $|S| + |T| + |T^S| = |S| + |T| + (|G| - |T + S|) = |G| + 1$.

- If $|T| \leq |T^S|$, then $|S| + |T| \leq \frac{2(|S| + |T| + |T^S|)}{3} = \frac{2|G| + 2}{3}$, and (iii) holds by Theorem 13.

- $|T| > |T^S|$, then $|S| + |T^S| \leq \frac{2(|S|+|T|+|T^S|)}{3} = \frac{2|G|+2}{3}$.

Assume first $|U| \geq |S|$. By Theorem 13, S a quasi-periodic modular H -progression, where H is the hyper-atom of S . By Lemma 11, T a quasi-periodic modular H -progression. Thus (iii) holds.

Assume now $|U| < |S|$ and that U generates G . By Theorem 13, S a quasi-periodic modular H -progression, where H is the hyper-atom of U . By Lemma 11, T a quasi-periodic modular H -progression. Thus (iii) holds.

- If $\langle U \rangle \neq G$, then by Lemma 10, S and T are $\langle U \rangle$ -quasi-periodic. Thus (ii) holds.

The proof is complete. ■

Following a suggestion of Kemperman [12], we formulate the main classical critical pair result in the following way:

Corollary 15 (*Kemperman Structure Theorem [12]*)

Let A, B be finite subsets of an abelian group G with $|G| \geq 2$.

Then the following conditions are equivalent:

- (I) $|A + B| = |A| + |B| - 1$, and moreover $|(c - A) \cap B| = 1$ for some c if $A + B$ is periodic.
- (II) There is a nonzero subgroup H and H -quasi-periodic decompositions $A = A_0 \cup \dots \cup A_u$ and $B = B_0 \cup \dots \cup B_t$ such that (A_0, B_0) is an elementary pair and $|(\phi(A_0 + B_0) - \phi(A)) \cap \phi(B)| = 1$, where $\phi : G \mapsto G/H$ is the canonical morphism.

Proof.

The implication (II) $\Rightarrow$ (I) is quite easy. Suppose that (I) holds. Without loss of generality we may take $|A| \geq |B|$, $0 \in A \cap B$ and $G = \langle A \cup B \rangle$. Let Q denotes the period of $A + B$ and let $\psi : G \mapsto G/Q$ denotes the canonical morphism.

Case 1. $|Q| = 1$.

Clearly (II) holds with $H = \{0\}$ if $\{A, B\}$ is a weak pair. Suppose that $\{A, B\}$ is not a weak pair. By Theorem 14, there is a proper subgroup H such that A and B are H -quasi-periodic. Take a minimal such a group. By $\rho_H(X)$ we shall mean the unique non-full H -coset's trace on X if such a coset exists.

The pair $\{\rho(A), \rho(B)\}$ must be a weak pair, by Theorem 14. Since $A + B$ is aperiodic, we have $|(\phi(\rho(A) + \rho(B)) - \phi(A)) \cap \phi(B)| = 1$, where $\phi : G \mapsto G/H$ is the canonical morphism.

Case 2. $|Q| \geq 2$.

Take two Q -decompositions $A = A_0 \cup \dots \cup A_u$ and $B = B_0 \cup \dots \cup B_t$ such that $c \in A_0 + B_0$. Observe that $\psi(c) = \psi(A_0) + \psi(B_0)$ has a unique expression. Hence by Scherck's Theorem C, $|\psi(A) + \psi(B)| \geq |\psi(A)| + |\psi(B)| - 1 = t + u + 1$. We must have $|\psi(A) + \psi(B)| = t + u + 1$, since otherwise $|A + B| = |\psi(A + B)||H| \geq |A| + |B|$.

By Lemma B, we have

$$|A_0| + |B_0| \leq |H| + 1.$$

Now we have

$$\begin{aligned} \sum_{0 \leq i \leq u} |A_i| + \sum_{0 \leq i \leq t} |B_i| - 1 &= |A + B| \\ &= |H|(u + t + 1) \geq |H|(u + t) + |A_0| + |B_0| - 1 \geq |A| + |B| - 1, \end{aligned}$$

observing that $|A| + |B| - 1 = |A + B| = |\psi(A + B)||H| = |A + H| + |B + H| - |H|$. It follows that

- $|A_1| = \cdots = |A_u| = |B_1| = \cdots = |B_t| = |H|$,
- $|A_0| + |B_0| \leq |H| + 1$.

Therefore $\{A_0, B_0\}$ is a weak pair. ■

We shall now prove Lev's Structure Theorem.

Corollary 16 (*Lev's Structure Theorem*)[14]

Let A and B be finite nonempty subsets of a non-zero abelian group G , with $|A| \leq |B|$, and $|A + B| = |A| + |B| - 1$. Suppose that either $A + B$ is aperiodic or there is a c with $|(c - A) \cap B| = 1$.

Then there exists a finite proper subgroup $H \subset G$ such that A and B are H -quasi-periodic and $\{\phi(A), \phi(B)\}$ is a weak pair.

Proof.

Without loss of generality we may assume that $0 \in A \cap B$ and $\langle A \cup B \rangle = G$. Put $L = \langle B \rangle$.

Let P be the period of $A + B$ and let $\sigma : G \mapsto G/P$ be the canonical morphism.

Case 1. $|P| = 1$.

- (i) If $G \neq L$ then A is L -quasi-periodic and $|\phi(B)| = 1$, where $\phi : G \mapsto G/L$ is the canonical morphism, by Theorem 14.
- (ii) If $G = L \neq \langle U \rangle$, then by Theorem 14, B and A are $\langle U \rangle$ -quasi-periodic, where $U = A^B - a$, for some $a \in A^B$. Also by Theorem 14, $\phi(B + A) = G/\langle U \rangle$, where $\phi : G \mapsto G/\langle U \rangle$ is the canonical morphism.
- (iii) If $G = L = \langle U \rangle$, then there is a proper subgroup H such that B and A are H -quasi-periodic modular progressions, by Theorem 14.

In order to show that $\{\phi(B), \phi(A)\}$ is a weak pair, it would be enough to observe that there is a uniquely representable element in $\phi(B) + \phi(A)$. This follows easily since $A + B$ is aperiodic and since A and B are quasi-periodic.

Case 2. $A + B$ is periodic.

By Case 1, there is subgroup K of G/P such that $\sigma(A)$ and $\sigma(B)$ are K -quasi-periodic and $\{\tau(\sigma(A)), \tau(\sigma(B))\}$ is a weak pair, where $\tau : G/K \mapsto G/P/K$. Put $\psi = \tau \circ \sigma$ and $Q = \psi^{-1}(K)$. Clearly A and B are Q -quasi-periodic. Clearly $\{\psi(A), \psi(B)\}$ is a weak pair. ■

References

- [1] E. Balandraud, Un nouveau point de vue isopérimétrique appliqué au théorème de Kneser, *Preprint*, december 2005.
- [2] D. Grynkiewicz, Quasi-periodic decompositions and the Kemperman’s structure theorem, *European J. Combin.* 26 (2005), no. 5, 559–575.
- [3] D. Grynkiewicz, A step beyond Kemperman’s structure Theorem, *Preprint* May 2006.
- [4] Y.O. Hamidoune, Sur les atomes d’un graphe orienté, *C.R. Acad. Sc. Paris A* 284 (1977), 1253–1256.
- [5] Y.O. Hamidoune, Quelques problèmes de connexité dans les graphes orientés, *J. Comb. Theory B* 30 (1981), 1–10.
- [6] Y.O. Hamidoune, On the connectivity of Cayley digraphs, *Europ. J. Combinatorics*, 5 (1984), 309–312.
- [7] Y.O. Hamidoune, Subsets with small sums in abelian groups I: The Vosper property. *European J. Combin.* 18 (1997), no. 5, 541–556.
- [8] Y.O. Hamidoune, An isoperimetric method in additive theory. *J. Algebra* 179 (1996), no. 2, 622–630.
- [9] Y.O. Hamidoune, Some results in Additive number Theory I: The critical pair Theory, *Acta Arith.* 96, no. 2(2000), 97–119.
- [10] Y.O. Hamidoune, Some additive applications of the isoperimetric approach, *Ann. Institut Fourier, Grenoble*, to appear, <http://arxiv.org/abs/math/07060635>.
- [11] Y. O. Hamidoune , A. Plagne. A new critical pair theorem applied to sum-free sets. *Comment. Math. Helv.* 79 (2004), no. 1, 183–207.
- [12] J. H. B. Kemperman, On small sumsets in Abelian groups, *Acta Math.* 103 (1960), 66–88.
- [13] R. A. LEE, Proving Kneser’s theorem for finite groups by another e -transform *Proc. Amer. Math. Soc.* 44 (1974), 255–258.
- [14] V. F. Lev, Critical pairs in abelian groups and Kemperman’s structure theorem. *Int. J. Number Theory* 2 (2006), no. 3, 379–396.
- [15] P. Scherk, L.Moser, Advanced Problems and Solutions: Solutions: 4466, *Amer. Math. Monthly* 62 (1955), no. 1, 46–47.
- [16] T. Tao, V.H. Vu, *Additive Combinatorics*, Cambridge Studies in Advanced Mathematics 105 (2006), Cambridge University Press.
- [17] G. Vosper, The critical pairs of subsets of a group of prime order, *J. London Math. Soc.* 31 (1956), 200–205.

Acknowledgement. The author wishes an anonymous referee for valuable comments on a preliminary draft related to this work.